

Noncommutative Gravity on Moyal plane: Second order correction

V. Radovanović

Divčibare
Septembar 2013

Introduction

AdS gauge theory on commutative spacetime

Noncommutative $SO(2, 3)_*$ symmetry

Action

Conclusion

Outline

Introduction

1. M. Dimitrijević, V. Radovanović and H. Štefančić, Phys. Rev D (2012),
2. M. Dimitrijević, V. Radovanović, in preparation,
3. M. Dimitrijević, V. Radovanović and I. Simonović, in preparation

Introduction

Quantum field theory encounters problems at high energy/small distances. Some modifications are needed. One possibility is noncommutativity among space time coordinates. It is given by

$$[x^\mu, x^\nu] = i\theta^{\mu\nu}(x) .$$

Canonical noncommutativity

$$[x^\mu, x^\nu] = i\theta^{\mu\nu} = \text{const.}$$

Different modes are constructed on canonical NC space time:

ϕ^4 , QED, standard model, SUSY models;

renormalizability, unitarity, . . .

Introduction

Generalization general relativity or some other gravity theory (like Poincare gauge theory) on NC spacetime is a difficult task.

Many attempts:

- Twist approach

1. P.Aschieri, C.Blohmann, M.Dimitrijević, F. Meyer, P.Schupp and J. Wess, CQG **22**, 3511-3522 (2005),
2. P.Aschieri, M. Dimitrijević, F.Meyer and J.Wess, CQG **23**, 1883-1912 (2006), [hep-th/0510059].

Introduction

- Sieberg-Witten approach
 1. Ali Chamseedine, PLB. **504** (2001) 33;PRD. **69** (2004) 024015
 - 2 M. A. Cardella and D. Zanon, CQG **20** (2003) L95
 3. P. Aschieri and L. Castellani, JHEP (0906) (2009) 086
 4. P. Aschieri and L. Castellani, arXiv:1111.4822, ArXiv: 12051911
 5. R. Banerjee, P. Mukherjee and S. Samanta, PRD **75**, (2007) 125020
 6. Yang-Gang Miao, Zhao Xue and Shao-Jun Zhang, PRD **83**, (2011) 024023

AdS gauge theory on commutative spacetime

Consider a gauge theory with $SO(2, 3)$ as a gauge group in $4D$ Minkowski spacetime.

$SO(2, 3)$ is the isometry group anti de Sitter space.

Anti de Sitter space is a maximally symmetric space with a negative constant curvature.

M_{AB} -generators of $SO(2, 3)$ group

A, B, \dots take values $0, 1, 2, 3, 5$.

Commutation relations:

$$[M_{AB}, M_{CD}] = i(\eta_{AD}M_{BC} + \eta_{BC}M_{AD} - \eta_{AC}M_{BD} - \eta_{BD}M_{AC}), \quad (1)$$

$\eta_{AB} = \text{diag}(+, -, -, -, +)$ is $5D$ metric.

AdS gauge theory on commutative spacetime

Clifford generators Γ_A in $5D$ Minkowski space satisfy

$$\{\Gamma_A, \Gamma_B\} = 2\eta_{AB} . \quad (2)$$

M_{AB} are

$$M_{AB} = \frac{i}{2}[\Gamma_A, \Gamma_B] . \quad (3)$$

γ_a , ($a = 0, 1, 2, 3$) are the gamma matrices in $4D$ Minkowski spacetime

The gamma matrices in $5D$ are

$$\Gamma_A = (i\gamma_a\gamma_5, \gamma_5) .$$

γ_5 is defined by $\gamma^5 = \gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$.

AdS gauge theory on commutative spacetime

It is easy to show that

$$\begin{aligned} M_{ab} &= \frac{i}{4} [\gamma_a, \gamma_b] = \frac{1}{2} \sigma_{ab} , \\ M_{5a} &= \frac{i}{2} \gamma_a . \end{aligned} \tag{4}$$

If we introduce momenta $P_a = \frac{1}{l} M_{a5}$, where l is a constant with dimensions of length AdS algebra (1) becomes

$$\begin{aligned} [M_{ab}, M_{cd}] &= i(\eta_{ad} M_{bc} + \eta_{bc} M_{ad} - \eta_{ac} M_{bd} - \eta_{bd} M_{ac}) \\ [M_{ab}, P_c] &= i(\eta_{bc} P_a - \eta_{ac} P_b) \\ [P_a, P_b] &= -i \frac{1}{l^2} M_{ab} . \end{aligned} \tag{5}$$

In the limit $l \rightarrow \infty$ AdS algebra reduces usual Poincare algebra in 4D spacetime. (Wigner-Inonu contraction)

AdS gauge theory on commutative spacetime

ψ spinor matter field in the fundamental representation

Under the infinitesimal $SO(2, 3)$ gauge transformations it transforms as

$$\delta_\epsilon \psi = i\epsilon \psi = \frac{i}{2} \epsilon^{AB} M_{AB} \psi \quad (6)$$

The covariant derivative in the fundamental representation

$$D_\mu \psi = \partial_\mu \psi - i\omega_\mu \psi, \quad (7)$$

$$\omega_\mu = \frac{1}{2} \omega_\mu^{AB} M_{AB} = \frac{1}{4} \omega_\mu^{ab} \sigma^{ab} - \frac{1}{2} \omega_\mu^{a5} \gamma_a \quad (8)$$

is the $SO(2, 3)$ gauge potential. Decomposition: ω_μ^{AB} to ω_μ^{ab} , ω_μ^{a5} ,
 ω_μ^{ab} is a spin connection

$\omega_\mu^{a5} = \frac{1}{l} e_\mu^a$ are vielbeins (tetrads).

AdS gauge theory on commutative spacetime

The transformation law of the $SO(2, 3)$ potential is given by

$$\delta_\epsilon \omega_\mu^{AB} = \partial_\mu \epsilon^{AB} - \epsilon^A_C \omega_\mu^{CB} + \epsilon^B_C \omega_\mu^{CA} . \quad (9)$$

If $\epsilon^{a5} = 0$ we obtain the transformation laws for spin connection and vielbeins under the $SO(1, 3)$ gauge transformation:

$$\delta_\epsilon \omega_\mu^{ab} = \partial_\mu \epsilon^{ab} - \epsilon^a_c \omega_\mu^{cb} + \epsilon^b_a \omega_\mu^{ca} , \quad (10)$$

$$\delta_\epsilon e_\mu^a = -\epsilon^{ad} e_\mu^d \quad (11)$$

We reduce the local anti de Sitter symmetry down to the local Lorentz symmetry:

$$SO(2, 3) \rightarrow SO(1, 3)$$

AdS gauge theory on commutative spacetime

The field strength

$$\begin{aligned}
 F_{\mu\nu} &= \partial_\mu \omega_\nu - \partial_\nu \omega_\mu - i[\omega_\mu, \omega_\nu] = \frac{1}{2} F_{\mu\nu}^{AB} M_{AB} \\
 &= \frac{1}{2} F_{\mu\nu}^{ab} M_{ab} + F_{\mu\nu}^{a5} M_{a5} ,
 \end{aligned} \tag{12}$$

where

$$F_{\mu\nu}^{ab} = R_{\mu\nu}^{ab} - \frac{1}{l^2} (e_\mu^a e_\nu^b - e_\mu^b e_\nu^a) . \tag{13}$$

Reiman curvature tensor is

$$R_{\mu\nu}^{ab} = \partial_\mu \omega_\nu^{ab} - \partial_\nu \omega_\mu^{ab} + \omega_\mu^{ac} \omega_\nu^{cb} - \omega_\mu^{bc} \omega_\nu^{ca} \tag{14}$$

Torsion

$$l F_{\mu\nu}^{a5} = D_\mu e_\nu^a - D_\nu e_\mu^a = T_{\mu\nu}^a \tag{15}$$

AdS gauge theory on commutative spacetime

Under the local anti de Sitter transformation field strength transforms as

$$\delta_\epsilon F_{\mu\nu} = i[\epsilon, F_{\mu\nu}] \quad (16)$$

or more explicitly

$$\begin{aligned} \delta_\epsilon F_{\mu\nu}^{ab} &= -\epsilon^{ac} F_{\mu\nu c}^b + \epsilon^{bc} F_{\mu\nu c}^a - \epsilon^{a5} F_{\mu\nu 5}^b + \epsilon^{b5} F_{\mu\nu 5}^a \\ \delta_\epsilon T_{\mu\nu}^a &= -\epsilon^{ac} T_{\mu\nu c}^a + \epsilon^{5c} F_{\mu\nu c}^a . \end{aligned} \quad (17)$$

If we reduce the initial anti de Sitter gauge symmetry down to its Lorentz subgroup by taking $\epsilon^{a5} = 0$ we obtain the correct transformation laws:

$$\begin{aligned} \delta_\epsilon F_{\mu\nu}^{ab} &= -\epsilon^a_c F_{\mu\nu}^{cb} + \epsilon^b_c F_{\mu\nu}^{ca} \\ \delta_\epsilon T_{\mu\nu}^a &= -\epsilon^a_c T_{\mu\nu}^c . \end{aligned} \quad (18)$$

AdS gauge theory on commutative spacetime

Action:

$$S = \frac{i l}{64\pi G_N} \int \epsilon^{\mu\nu\rho\sigma} \text{Tr} (F_{\mu\nu} F_{\rho\sigma} \phi) + \lambda \int d^4x \left(\frac{1}{4} \text{Tr} \phi^2 - l^2 \right) \quad (19)$$

is invariant under the $SO(2, 3)$ gauge transformations

G_N is the Newton gravitational constant.

λ is the Lagrange multiplier.

$\phi = \phi^A \Gamma_A$ is a vector field

$$\delta\phi = i[\epsilon, \phi] , \quad (20)$$

Constraint $\phi_A \phi^A = l^2$.

AdS gauge theory on commutative spacetime

Taking $\phi^a = 0$, $\phi^5 = l$ the $SO(2, 3)$ symmetry is broken spontaneously down to $SO(1, 3)$

The action after SSB:

$$S = \frac{1}{16\pi G_N} \int d^4x \left[\frac{l^2}{16} \epsilon^{\mu\nu\rho\sigma} \epsilon_{abcd} R_{\mu\nu}^{ab} R_{\rho\sigma}^{cd} + \sqrt{-g}R - 2\sqrt{-g}\Lambda \right], \quad (21)$$

where $\Lambda = -3/l^2$ and $\sqrt{-g} = \det e_{\mu}^a$.

AdS gauge theory on commutative spacetime

In this action (see the third line) the vielbeins and spin connection are independent variables. Varying the action with respect to the spin connection we obtain an equation which relates connection and vielbeins. In this way we can express the spin connection in terms of vielbeins. Since there is no fermionic matter in the action (21) this equation gives the vanishing torsion. In that case the first term in (21) is the Gauss-Bonnet term; it is a topological term and does not contribute to the equations of motion. The second term is the Einstein-Hilbert action, while the last term is the cosmological constant.

AdS gauge theory on commutative spacetime

From vielbeins we can construct the metric tensor

$$g_{\mu\nu} = \eta_{ab} e_{\mu}^a e_{\nu}^b . \quad (22)$$

Metric tensor can be constuced from field Φ by

$$g_{\mu\nu} = \eta_{AB} D_{\mu} \Phi^A D_{\nu} \Phi^B . \quad (23)$$

The action (21) is invariant under the Lorentz gauge transformations by construction. In addition this action posses invariance under general coordinate transformations. This action will be our starting point for the construction of a noncommutative gravity theory.

References:

MacDowell, Mansurri, PRL (1977)

Stelle, West, PRD, 1980

P. Townsend, PRD, 1977

AdS gauge theory on commutative spacetime

Interes za ovaj pristup gravitaciji raste:

1. A. Chamseddine and V. Mukhanov, 'Gravity with de Sitter and unitary tangent groups' JHEP (2010)
2. J. Barret and S. Kerr, 'Gauge gravity and discrete quantum models' Arxiv: 1309.1660

Noncommutative $SO(2, 3)_*$ symmetry

Replace the usual product by the Moyal-Weyl \star product

$$f(x) \star g(x) = e^{\frac{i}{2} \theta^{\mu\nu} \frac{\partial}{\partial x^\mu} \frac{\partial}{\partial y^\nu}} f(x)g(y)|_{y \rightarrow x} , \quad (24)$$

where $\theta^{\mu\nu}$ is a constant antisymmetric matrix,

Replace the commutative fields by their noncommutative counterparts.

Noncommutative gauge potential $\hat{\omega}_\mu$

Noncommutative curvature tensor $\hat{F}_{\mu\nu}$

$$\hat{F}_{\mu\nu} = \partial_\mu \hat{\omega}_\nu - \partial_\nu \hat{\omega}_\mu - i[\hat{\omega}_\mu \star \hat{\omega}_\nu] . \quad (25)$$

Noncommutative adjoint field $\hat{\Phi}$

Noncommutative $SO(2,3)_*$ symmetry

Under the deformed gauge transformations the gauge potential, the field strength and $\hat{\Phi}$ field transform as

$$\begin{aligned}\delta_\epsilon^* \hat{\omega}_\mu &= \partial_\mu \hat{\Lambda}_\epsilon - i[\hat{\omega}_\mu \star \hat{\Lambda}_\epsilon] \\ \delta_\epsilon^* \hat{F}_{\mu\nu} &= i[\hat{\Lambda}_\epsilon \star \hat{F}_{\mu\nu}],\end{aligned}\tag{26}$$

$$\delta_\epsilon^* \hat{\Phi} = i[\hat{\Lambda}_\epsilon \star \hat{F}]\tag{27}$$

where $\hat{\Lambda}_\epsilon$ is the noncommutative gauge parameter.

The noncommutative fields belong to the enveloping algebra of $so(2,3)$. For example, the \star -commutator in (??) does not close in the Lie algebra.

Noncommutative $SO(2, 3)_*$ symmetry

Commutative and noncommutative symmetries which correspond to the same gauge group can be related by the Seiberg-Witten map: the map enables one to express the noncommutative variables in terms of the commutative variables. In that way no new degrees of freedom are introduced. SW map can also be seen as an expansion in $\theta^{\mu\nu}$, so the SW approach is known as a θ -expanded theory.

Noncommutative $SO(2, 3)_*$ symmetry

The noncommutative quantities $\hat{\Lambda}_\epsilon, \hat{\omega}_\mu, \hat{\Phi}$ are power series in the noncommutative parameter $\theta^{\mu\nu}$:

$$\begin{aligned}\hat{\Lambda}_\epsilon &= \epsilon + \hat{\Lambda}^{(1)} + \hat{\Lambda}^{(2)} + \dots, \\ \hat{\omega}_\mu &= \omega_\mu + \hat{\omega}_\mu^{(1)} + \hat{\omega}_\mu^{(2)} + \dots,\end{aligned}\tag{28}$$

where the higher order corrections are functions of the commutative variables ϵ, ω_μ , and their derivatives. The requirement that the commutator of two deformed gauge transformations is a deformed transformation again:

$$[\delta_\alpha^* ; \delta_\beta^*] = \delta_{-i[\alpha, \beta]}^*\tag{29}$$

gives the solution for $\Lambda_\epsilon^{(1)}, \Lambda_\epsilon^{(2)}, \dots$

Noncommutative $SO(2, 3)_*$ symmetry

Solving the equation

$$\hat{\omega}_\mu(\omega) + \delta_\epsilon^* \hat{\omega}_\mu(\omega) = \hat{\omega}_\mu(\omega + \delta_\epsilon \omega) \quad (30)$$

order by order in the noncommutative parameter we can express noncommutative gauge potential $\hat{\omega}_\mu$ in terms of the commutative one. The first order solution is

$$\hat{\omega}_\mu^{(1)} = -\frac{1}{4} \theta^{\kappa\lambda} \{ \omega_\kappa, \partial_\lambda \omega_\mu + F_{\lambda\mu} \} \quad (31)$$

Noncommutative $SO(2, 3)_*$ symmetry

From potential $\hat{\omega}_\mu$ we find the field strength

$$\hat{F}_{\mu\nu} = \partial_\mu \hat{\omega}_\nu - \partial_\nu \hat{\omega}_\mu - i[\hat{\omega}_\mu * \hat{\omega}_\nu], \quad (32)$$

The first order correction is

$$\hat{F}_{\mu\nu}^{(1)} = -\frac{1}{4}\theta^{\kappa\lambda}\{\omega_\kappa, \partial_\lambda F_{\mu\nu} + D_\lambda F_{\mu\nu}\} + \frac{1}{2}\theta^{\kappa\lambda}\{F_{\mu\kappa}, F_{\nu\lambda}\} \quad (33)$$

Transformation law

$$\delta_\epsilon^* \hat{F}_{\mu\nu} = i[\hat{\Lambda}_\epsilon * \hat{F}_{\mu\nu}] \quad (34)$$

Action

The NC action is given by

$$S_{NC} = -\frac{i l}{16\pi G_N} \text{Tr} \int d^4x \epsilon^{\mu\nu\rho\sigma} \hat{F}_{\mu\nu} \star \hat{F}_{\rho\sigma} \star \hat{\phi}. \quad (35)$$

The \star -product is the Moyal-Weyl \star -product;
fields with "hat" are NC fields;

The action is invariant under the NC $SO(2, 3)$ gauge group.

Action

Seiberg-Witten expansion

$$S_{NC} = S^{(0)} + S^{(1)} + S^{(2)} + \dots \quad (36)$$

Under NC gauge transformation the 'field' $(\hat{F}_{\alpha\beta} \star \hat{F}_{\mu\nu})^{(1)}$ transforms in adjoint representation

$$\begin{aligned} (\hat{F}_{\alpha\beta} \star \hat{F}_{\mu\nu})^{(1)} &= F_{\mu\nu}^{(1)} F_{\rho\sigma} + F_{\mu\nu} F_{\rho\sigma}^{(1)} + \frac{i}{2} \theta^{\alpha\beta} \partial_\alpha F_{\mu\nu} \partial_\beta F_{\rho\sigma} \\ &= -\frac{1}{4} \theta^{\kappa\lambda} \{ \omega_\kappa, \partial_\lambda (F_{\alpha\beta} F_{\mu\nu}) + D_\lambda (F_{\alpha\beta} F_{\mu\nu}) \} \\ &\quad + \frac{i}{2} (D_\kappa F_{\alpha\beta}) (D_\lambda F_{\mu\nu}) + \frac{i}{2} \theta^{\kappa\lambda} (\{ F_{\kappa\alpha}, F_{\lambda\beta} \} F_{\mu\nu} \\ &\quad + F_{\alpha\beta} \{ F_{\kappa\mu}, F_{\lambda\nu} \}) \end{aligned}$$

Action

$$\begin{aligned}
 & (\hat{F}_{\alpha\beta} \star \hat{F}_{\mu\nu} \star \hat{\phi})^{(1)} \\
 = & (F_{\alpha\beta} \star F_{\mu\nu})^{(1)} \phi + (F_{\alpha\beta} F_{\mu\nu}) \phi^{(1)} + \frac{i}{2} \partial_\alpha (F_{\alpha\beta} F_{\mu\nu}) \partial_\beta \phi \\
 = & -\frac{1}{4} \theta^{\alpha\beta} \{ \omega_\alpha, (\partial_\beta + D_\beta) (F_{\alpha\beta} F_{\mu\nu} \phi) \} + \frac{i}{2} \theta^{\alpha\beta} D_\alpha (F_{\mu\nu} F_{\rho\sigma}) D_\beta \phi \\
 + & \frac{i}{2} \theta^{\alpha\beta} D_\alpha F_{\mu\nu} D_\beta F_{\rho\sigma} \phi \\
 + & \frac{1}{2} \theta^{\alpha\beta} \{ F_{\alpha\mu}, F_{\beta\nu} \} F_{\rho\sigma} \phi + \frac{1}{2} \theta^{\alpha\beta} F_{\mu\nu} \{ F_{\alpha\rho}, F_{\beta\sigma} \} \phi \quad (37)
 \end{aligned}$$

Action

The NC action in first order in θ is

$$\begin{aligned}
 S_{NC}^{(1)} &= -\frac{i\ell}{16\pi G_N} \text{Tr} \int d^4x \epsilon^{\mu\nu\rho\sigma} (\hat{F}_{\mu\nu} \star \hat{F}_{\rho\sigma} \star \hat{\phi})^{(1)} \\
 &= -\frac{i\ell}{16\pi G_N} \theta^{\alpha\beta} \text{Tr} \int d^4x \epsilon^{\mu\nu\rho\sigma} \left(-\frac{1}{4} F_{\mu\nu} F_{\rho\sigma} \{F_{\alpha\beta}, \phi\} \right. \\
 &\quad + \frac{i}{2} D_\alpha F_{\mu\nu} D_\beta F_{\rho\sigma} \phi \\
 &\quad \left. + \frac{1}{2} \{F_{\alpha\mu}, F_{\beta\nu}\} F_{\rho\sigma} \phi + \frac{1}{2} F_{\mu\nu} \{F_{\alpha\rho}, F_{\beta\sigma}\} \phi \right) \quad (38)
 \end{aligned}$$

Result $S_{NC}^{(1)} = 0$.

Action

From the first order we can easily find the second order.

$$\begin{aligned}
 S_{NC}^{(2)} &= -\frac{i l}{128\pi G_N} \theta^{\alpha\beta} \text{Tr} \int d^4x \epsilon^{\mu\nu\rho\sigma} \left(-\frac{1}{4} \hat{F}_{\mu\nu} \star \hat{F}_{\rho\sigma} \star \{ \hat{F}_{\alpha\beta} \star \hat{\phi} \} \right. \\
 &+ \frac{i}{2} D_\alpha \hat{F}_{\mu\nu} \star D_\beta \hat{F}_{\rho\sigma} \star \phi \\
 &+ \left. \frac{1}{2} \{ \hat{F}_{\alpha\mu} \star \hat{F}_{\beta\nu} \} \star \hat{F}_{\rho\sigma} \star \hat{\phi} + \frac{1}{2} \hat{F}_{\mu\nu} \star \{ \hat{F}_{\alpha\rho} \star \hat{F}_{\beta\sigma} \} \star \hat{\phi} \right)^{(1)}
 \end{aligned}$$

Action

$$\begin{aligned}
S_{NC}^{(2)} = & \frac{i l}{256 \pi G_N} \theta^{\alpha \beta} \theta^{\kappa \lambda} \text{Tr} \int d^4 x \epsilon^{\mu \nu \rho \sigma} \left(-\frac{1}{16} \{F_{\alpha \beta}, \{F_{\mu \nu}, F_{\rho \sigma}\}\} \{\phi, F_{\kappa \lambda}\} \right. \\
& + \frac{1}{4} \{F_{\alpha \beta}, \{F_{\rho \sigma}, \{F_{\kappa \mu}, F_{\lambda \nu}\}\}\} \phi + \frac{1}{8} \{ \{F_{\mu \nu}, F_{\rho \sigma}\}, \{F_{\kappa \alpha}, F_{\lambda \beta}\} \} \phi \\
& + \frac{i}{8} \{F_{\alpha \beta}, [D_{\kappa} F_{\mu \nu}, D_{\lambda} F_{\rho \sigma}]\} \phi + \frac{i}{4} [\{D_{\kappa} F_{\mu \nu}, F_{\rho \sigma}, D_{\lambda} F_{\alpha \beta}\}] \phi \\
& - \frac{1}{4} \{F_{\rho \sigma}, \{F_{\alpha \mu}, F_{\beta \nu}\}\} \{\phi, F_{\kappa \lambda}\} + \frac{1}{2} \{ \{F_{\alpha \mu}, F_{\beta \nu}, \{F_{\kappa \rho}, F_{\lambda \sigma}\}\} \} \phi \\
& + \{F_{\rho \sigma}, \{F_{\beta \nu}, \{F_{\kappa \alpha}, F_{\lambda \mu}\}\}\} \phi + \frac{i}{2} \{F_{\rho \sigma}, [D_{\kappa} F_{\alpha \mu}, D_{\lambda} F_{\beta \nu}]\} \phi \\
& + [\{F_{\beta \nu}, D_{\kappa} F_{\alpha \mu}\}, D_{\lambda} F_{\rho \sigma}] \phi \\
& - \frac{i}{8} \{ \phi, F_{\kappa \lambda} \} [D_{\alpha} F_{\mu \nu}, D_{\beta} F_{\rho \sigma}] - \frac{1}{4} \{D_{\kappa} D_{\alpha} F_{\mu \nu}, D_{\lambda} D_{\beta} F_{\rho \sigma}\} \phi \\
& + \frac{i}{8} [\{F_{\alpha \beta}, D_{\kappa} F_{\mu \nu}\}, D_{\lambda} F_{\rho \sigma}] + \frac{i}{8} [\{F_{\alpha \beta}, D_{\lambda} F_{\rho \sigma}\}, D_{\kappa} F_{\mu \nu}]
\end{aligned}$$

Action

After the symmetry breaking the field $\phi^a = 0$, $\phi^5 = l$.

$$\begin{aligned}
 S_{NC}^{(2)} = & -\frac{l^2}{64\pi G_N} \theta^{\alpha\beta} \theta^{\kappa\lambda} \epsilon^{\mu\nu\rho\sigma} \epsilon_{abcd} \int d^4x \left[\frac{1}{256} \left(F_{\mu\nu}{}^{cd} F_{\rho\sigma}{}^{ab} F_{\alpha\beta}{}^{mn} F_{\kappa\lambda mn} \right. \right. \\
 - & \left. \left. 8F_{\mu\nu}{}^{ab} F_{\rho\sigma}{}^{c5} F_{\kappa\lambda}{}^{de} F_{\alpha\beta e}{}^5 + F_{\alpha\beta}{}^{ab} F_{\kappa\lambda}{}^{cd} (F_{\mu\nu}{}^{mn} F_{\rho\sigma mn} + 2F_{\mu\nu}{}^{m5} F_{\rho\sigma m}{}^{5b}) \right) \right. \\
 - & \left. \frac{1}{32} \left(F_{\kappa\lambda}{}^{ab} F_{\mu\nu}{}^{cd} F_{\alpha\rho}{}^{mn} F_{\beta\sigma mn} + 2F_{\alpha\beta}{}^{ab} F_{\rho\sigma}{}^{cd} F_{\kappa\mu}{}^{m5} F_{\lambda\nu m}{}^5 + F_{\kappa\mu}{}^{ab} F_{\lambda\nu}{}^{cd} F_{\alpha\beta}{}^{mn} F_{\rho} \right. \right. \\
 - & \left. \left. \frac{1}{128} \left(F_{\kappa\alpha}{}^{ab} F_{\lambda\beta}{}^{cd} (F_{\mu\nu}{}^{mn} F_{\rho\sigma mn} + 2F_{\mu\nu}{}^{m5} F_{\rho\sigma m}{}^5) \right) \right. \right. \\
 + & \left. \left. F_{\mu\nu}{}^{ab} F_{\rho\sigma}{}^{cd} (F_{\kappa\alpha}{}^{mn} F_{\lambda\beta mn} + 2F_{\kappa\alpha}{}^{m5} F_{\lambda\beta m}{}^5) \right) \right. \\
 + & \left. \frac{1}{16} F_{\alpha\beta}{}^{ab} \left((D_\kappa F_{\mu\nu})^{cm} (D_\lambda F_{\rho\sigma})_m^d + (D_\kappa F_{\mu\nu})^{c5} (D_\lambda F_{\rho\sigma})_5^d \right) \right]
 \end{aligned}$$

Action

$$\begin{aligned}
 & - \frac{1}{16} \left((D_\kappa F_{\mu\nu})^{ab} (D_\lambda F_{\alpha\beta})^{d5} F_{\rho\sigma}^{c5} + (D_\kappa F_{\mu\nu})^{a5} (D_\lambda F_{\alpha\beta})^{b5} F_{\rho\sigma}^{cd} \right) \\
 & + \frac{1}{16} F_{\alpha\mu}^{ab} F_{\beta\nu}^{cd} \left(F_{\kappa\rho}{}^{mn} F_{\lambda\sigma mn} + 2F_{\kappa\rho}{}^{m5} F_{\lambda\sigma m5} \right) \\
 & + \frac{1}{16} \left(F_{\rho\sigma}{}^{ab} F_{\beta\nu}{}^{cd} (F_{\kappa\alpha}{}^{mn} F_{\lambda\mu mn} + 2F_{\kappa\alpha}{}^{m5} F_{\lambda\mu m5}) + F_{\kappa\alpha}{}^{ab} F_{\lambda\mu}{}^{cd} F_{\rho\sigma}{}^{mn} F_{\beta\nu mn} \right. \\
 & \left. - 4(F_{\kappa\alpha}{}^{ab} F_{\lambda\mu}{}^{c5} + F_{\kappa\alpha}{}^{a5} F_{\lambda\mu}{}^{bc}) F_{\rho\sigma}{}^{de} F_{\beta\nu e5} \right) \\
 & - \frac{1}{8} F_{\rho\sigma}{}^{ab} \left((D_\kappa F_{\alpha\mu})^{cm} (D_\lambda F_{\beta\nu})_m^d + (D_\kappa F_{\alpha\mu})^{c5} (D_\lambda F_{\beta\nu})_5^d \right) \\
 & + \frac{1}{2} \left(F_{\kappa\mu}{}^{ab} (D_\alpha F_{\lambda\nu})^{c5} (D_\beta F_{\rho\sigma})^{d5} + F_{\kappa\mu}{}^{a5} (D_\alpha F_{\lambda\nu})^{bc} \right) (D_\beta F_{\rho\sigma})^{d5} \\
 & + \frac{1}{8} \left(F_{\kappa\alpha}{}^{ab} (D_\lambda F_{\mu\nu})^{c5} + F_{\kappa\alpha}{}^{a5} (D_\lambda F_{\mu\nu})^{bc} \right) (D_\beta F_{\rho\sigma})^{d5} \\
 & - \frac{1}{32} (D_\kappa D_\alpha F_{\mu\nu})^{ab} (D_\lambda D_\beta F_{\rho\sigma})^{cd} \Big] \tag{41}
 \end{aligned}$$

Action

Terms in the action ($T = 0$)

$$\frac{\theta\theta}{l^4 G_N} \left(l^6 R^4, l^4 R^3, l^2 R^2, R, \frac{1}{l^2} \right) \quad (42)$$

Limesi:

1. $R \ll \frac{1}{l^2}$, , big cosmological constant, smaller energy
 2. $R \gg \frac{1}{l^2}$, small cosmological constant, higher energy
- $\Lambda_{\text{eff}} = (10^{-3} \text{eV})^4$

Action

$$\begin{aligned}
S^{(2)} = & \frac{3\theta^{\alpha\beta}\theta^{\kappa\lambda}}{64\pi G_N l^6} \int d^4x \sqrt{-g} g_{\alpha\kappa} g_{\beta\lambda} \\
& - \frac{\theta^{\alpha\beta}\theta^{\kappa\lambda}}{64\pi G_N l^4} \int d^4x \sqrt{-g} \left(3g_{\alpha\kappa} R_{\beta\lambda} + 3R_{\alpha\beta\kappa\lambda} - 2R_{\alpha\kappa\beta\lambda} \right) \\
& - \frac{\theta^{\alpha\beta}\theta^{\kappa\lambda}}{256\pi G_N l^2} \int d^4x \sqrt{-g} \left(2g_{\alpha\kappa} (-3RR_{\beta\lambda} + 4R_{\beta\mu} R_{\lambda}^{\mu} + 2R^{\mu\nu} R_{\beta\nu\lambda\mu} \right. \\
& \left. - 2R_{\beta\mu}{}^{\rho\sigma} R_{\rho\sigma\lambda}{}^{\mu}) \right. \\
& + R(2R_{\alpha\kappa\beta\lambda} - R_{\alpha\beta\kappa\lambda}) - 8R_{\alpha\kappa} R_{\beta\lambda} - 14R_{\alpha\beta\kappa\mu} R_{\lambda}^{\mu} - 16R_{\alpha\kappa\beta\mu} R_{\lambda}^{\mu} \quad (43) \\
& \left. - 2R_{\alpha\beta}{}^{\mu\nu} (2R_{\kappa\lambda\mu\nu} + 2R_{\kappa\mu\lambda\nu}) \right. \\
& \left. + 2R_{\alpha\kappa}{}^{\mu\nu} (5R_{\beta\lambda\mu\nu} - 4R_{\beta\mu\lambda\nu}) + 4R_{\alpha\mu\beta}{}^{\nu} R_{\kappa\nu\lambda}{}^{\mu} + 4R_{\alpha\mu\kappa}{}^{\nu} R_{\lambda\nu\beta}{}^{\mu} \right) \\
+ & \frac{-l^2\theta^{\alpha\beta}\theta^{\kappa\lambda}}{64\pi G_N 16} \int d^4x \epsilon^{\mu\nu\rho\sigma} \epsilon_{abcd} \left(-\frac{8}{l^6} e_{\alpha}^a e_{\beta}^b \nabla_{\kappa} (e_{\mu}^c e_{\nu}^m) \nabla_{\lambda} (e_{\rho m} e_{\sigma}^d) + \dots \right)
\end{aligned}$$

Action

$$\nabla_\mu e_\nu^a = \partial_\mu e_\nu^a + \omega_\mu^{ab} e_{\nu b} \quad (44)$$

is $SO(1, 3)$ covariant derivative. Comments:

$$\Lambda \rightarrow \Lambda_{\text{eff}}(x) = \Lambda + \frac{\theta^{\alpha\beta} \theta^{\kappa\lambda} g_{\alpha\kappa} g_{\beta\lambda}}{72} \Lambda^3 \quad (45)$$

Modification of Freedman solution?,
 mass for graviton?
 diffeomorphisms symmetry.

Action

On higher energy the higher powers in R are important.

$$S = \frac{l^2}{64\pi G_N} \theta^{\alpha\beta} \theta^{\kappa\lambda} \int d^4x \sqrt{-g} \left(\frac{1}{16} R_{\mu\nu}^{\gamma\delta} R_{\rho\sigma\gamma\delta} R_{\kappa\alpha}^{\rho\sigma} R_{\lambda\beta}^{\mu\nu} + \dots \right) \quad (46)$$

These terms might be important in early inflation expansion of Universe.

Action

We can start from commutative action

$$S = \frac{1}{64\pi G_N l} \int d^4x \epsilon^{\mu\nu\rho\sigma} \text{Tr} (F_{\mu\nu} D_\rho \Phi D_\sigma \Phi) \quad (47)$$

After SSB we get

$$S = -\frac{1}{16\pi G_N l} \int d^4x e (R - \frac{12}{l^2}) . \quad (48)$$

Deformation of this term is in progress (with I. Simonovic, M. Dimitrijevic)

1. H. Pagels, Phys. Rev D (1984)
2. F. Wilczek, Phys. Rev. Lett (1998)

Action

$$\begin{aligned}
 S &= \epsilon^{\mu\nu\rho\sigma} \int \text{Tr} ([F_{\mu\nu}, \Phi][F_{\rho\sigma}, \Phi]) \\
 &\sim \epsilon^{\mu\nu\rho\sigma} \int F_{\mu\nu}^{AB} \phi_B F_{\rho\sigma AC} \phi^C
 \end{aligned} \tag{49}$$

After SSB we get the action

$$S \sim \epsilon^{\mu\nu\rho\sigma} \int T_{\mu\nu}^a T_{\rho\sigma}^b \eta_{ab} . \tag{50}$$

Conclusion

We consider a deformation of gravity with local $SO(2, 3)_*$ symmetry; Breaking the commutative symmetry

$$SO(2, 3) \rightarrow SO(1, 3) \quad (51)$$

we get EH term, cosmological constant term and second order corrections in θ .

The second order corrections to the classical action are found in the covariant form.

Future investigations:

NC corrections to the black hole/ cosmological solutions

NC terms are important in early evolution of universe (inflationary expansion)

renormalizability of NC gravity